

1. 다음 중 계산 한 값이 옳은 것은?

$$\textcircled{1} \quad \sqrt{3^2} - \sqrt{(-5)^2} + \sqrt{2^2} = 10$$

$$\textcircled{2} \quad \sqrt{(-2)^2} - (-\sqrt{3})^2 - \sqrt{5^2} = 0$$

$$\textcircled{3} \quad \sqrt{\left(\frac{2}{5}\right)^2} + \sqrt{\frac{9}{25}} - \sqrt{\left(\frac{6}{5}\right)^2} = -\frac{1}{5}$$

$$\textcircled{4} \quad \sqrt{2^2} \times \sqrt{\left(\frac{1}{2}\right)^2} + \sqrt{\left(-\frac{1}{2}\right)^2} = 0$$

$$\textcircled{5} \quad \sqrt{3^2} + \sqrt{4^2} - \sqrt{(-5)^2} = 12$$

해설

$$\textcircled{1} \quad \sqrt{3^2} - \sqrt{(-5)^2} + \sqrt{2^2} = 3 - 5 + 2 = 0$$

$$\textcircled{2} \quad \sqrt{(-2)^2} - (-\sqrt{3})^2 - \sqrt{5^2} = 2 - 3 - 5 = -6$$

$$\textcircled{3} \quad \sqrt{\left(\frac{2}{5}\right)^2} + \sqrt{\frac{9}{25}} - \sqrt{\left(\frac{6}{5}\right)^2} = \frac{2}{5} + \frac{3}{5} - \frac{6}{5} = -\frac{1}{5}$$

$$\textcircled{4} \quad \sqrt{2^2} \times \sqrt{\left(\frac{1}{2}\right)^2} + \sqrt{\left(-\frac{1}{2}\right)^2} = 2 \times \frac{1}{2} + \frac{1}{2} = \frac{3}{2}$$

$$\textcircled{5} \quad \sqrt{3^2} + \sqrt{4^2} - \sqrt{(-5)^2} = 3 + 4 - 5 = 2$$

2.  $x > 2$  일 때, 다음 중  $\sqrt{(x-2)^2} - \sqrt{(2-x)^2}$  의 값은?

- ① -2      ② -1      ③ 0      ④ 1      ⑤ 2

해설

$x > 2$  ⇒  $x-2 > 0, 2-x < 0$

$$\begin{aligned}(\text{준식}) &= (x-2) - \{-(2-x)\} \\&= (x-2) - (x-2) = 0\end{aligned}$$

3. 다음 중 부등식  $4 < \sqrt{x} \leq 5$  를 만족하는 자연수  $x$  가 아닌 것은?

- ① 18      ② 20      ③ 22      ④ 24      ⑤ 26

해설

$$4 = \sqrt{16} < \sqrt{x} \leq 5 = \sqrt{25}$$

$$\therefore x = 17, 18, 19, 20, 21, 22, 23, 24, 25$$