

1. a 의 값의 범위가 $-2 < a < 2$ 일 때, $\sqrt{(a-2)^2} - \sqrt{(a+2)^2}$ 의 식을 간단히 하면?

① 0

② $-2a - 4$

③ -4

④ $-2a$

⑤ $2a$

2. $3 < x < 4$ 일 때, $\sqrt{(3-x)^2} - \sqrt{(x-4)^2}$ 을 간단히 하면?

① $2x - 1$

② $2x - 3$

③ $2x - 5$

④ $2x - 7$

⑤ $2x - 9$

3. $a > 0$ 일 때, $-\sqrt{(-5a)^2} + \sqrt{16a^2}$ 을 간단히 하여라.

 답: _____

4. $a > 0$ 일 때, $\sqrt{(-4a)^2} - \sqrt{9a^2} + (-\sqrt{2a})^2$ 을 간단히 하면?

① $-a$

② $3a$

③ $5a$

④ a

⑤ $-3a$

5. $a < 0$, $b > 0$ 일 때, $-\sqrt{b^2} - \sqrt{a^2}$ 을 간단히 하면?

① $b - a$

② $a - b$

③ $-a - b$

④ $a + b$

⑤ $-a^2 + b^2$

6. $1 < x < 3$ 일 때, $\sqrt{(x-3)^2} + \sqrt{(x+1)^2}$ 을 간단히 하여라.

 답: _____

7. $a > 0$ 일 때, 다음 계산에서 옳지 않은 것을 모두 골라라.

- $\textcircled{\tiny \neg} \quad -\sqrt{121a^2} - \sqrt{(-7a)^2} = -4a$
- $\textcircled{\tiny \text{L}} \quad \sqrt{25a^2} + \sqrt{(-6a)^2} = -a$
- $\textcircled{\tiny \text{E}} \quad -\sqrt{(-4a)^2} \times \frac{\sqrt{25a^2}}{a^2} = -20a$
- $\textcircled{\tiny \text{E}} \quad 9\sqrt{a^2} + \sqrt{(-6a)^2} - \sqrt{a^2} = 14a$

 답: _____

 답: _____

 답: _____

8. $a > 0, b < 0$ 일 때, $\sqrt{a^2} + \sqrt{(-b)^2} - \sqrt{4a^2} - \sqrt{b^2}$ 을 간단히 하면?

① $-a - b$

② $-a - 2b$

③ a

④ $-a$

⑤ $-a + 2b$

9. $a > 0$, $b < 0$ 일 때, $\sqrt{(2a)^2} + \sqrt{(-a)^2} - \sqrt{(5b)^2}$ 을 간단히 하면?

① $a - 5b$

② $a + 5b$

③ $3a - 5b$

④ $3a + 5b$

⑤ $5a - 5b$

10. $x < 0$ 일 때, $\sqrt{(-3x)^2} - \sqrt{(5x)^2} - \sqrt{(9x^2)}$ 을 간단히 하면?

- ① $-5x$ ② x ③ $5x$ ④ $11x$ ⑤ $13x$

11. $a > 0, b < 0$ 일 때, 다음 중 옳은 것은?

① $\sqrt{9a^2} \times \sqrt{(-6b)^2} = 18ab$

② $-\sqrt{(3a)^2} \div \sqrt{(-2a)^2} = \frac{3}{2}a$

③ $\sqrt{(-5b)^2} - \sqrt{16b^2} = b$

④ $2\sqrt{a^2} \times \sqrt{(-b)^2} + \sqrt{25a^2b^2} = -7ab$

⑤ $\sqrt{a^2} + \sqrt{(-b)^2} = a + b$

12. $0 < a < 1$ 일 때, $\sqrt{(a-1)^2} - \sqrt{(1-a)^2}$ 를 간단히 하라.

 답: _____

13. $0 < a < 1$ 일 때, $\sqrt{(1-a)^2} - \sqrt{(a-1)^2}$ 을 간단히 하면?

① 0

② 2

③ $2a - 2$

④ $2a + 2$

⑤ $-2a + 2$

14. $0 < x$ 일 때, $\sqrt{x^2} + \sqrt{(x+3)^2}$ 를 간단히 하면?

① 3

② $x + 3$

③ $x - 3$

④ $2x$

⑤ $2x + 3$

15. $a < 0$ 일 때, $\sqrt{81a^2} \div (-\sqrt{3a})^2 + \sqrt{(-0.5a)^2} \times \left(\sqrt{\frac{1}{5}a}\right)^2$ 을 계산하면?

① $0.1a^2 - 3$

② $0.1a^2 + 3$

③ $0.5a^2 - 3$

④ $0.5a^2 + 3$

⑤ $a^2 - 3$

16. 실수 a, b 에 대하여 $a < 0, ab < 0$ 일 때, $\sqrt{(2a-b)^2} + \sqrt{a^2} - \sqrt{(b-a)^2}$ 을 간단히 하면?

① $-4a + 2b$

② $-2a - 2b$

③ $-2a + 2b$

④ $-2a$

⑤ $4a - 2b$

17. $b < 0 < a < 2$ 일 때, 다음 중 옳은 것은?

① $\sqrt{(a-2)^2} = a-2$

② $\sqrt{(2-a)^2} = a-2$

③ $\sqrt{(a-b)^2} + \sqrt{(b-a)^2} = 0$

④ $\sqrt{b^2} + |b| = -2b$

⑤ $\sqrt{(b-2)^2} = b-2$

18. $-2 < x < 3$ 일 때, $\sqrt{(x+2)^2} - \sqrt{(x-3)^2} + 2|3-x|$ 를 간단히 하여라.

 답: _____

19. $a < 0$ 일 때, $A = \sqrt{(-3a)^2} \times (-\sqrt{a})^2 \div \sqrt{4a^2} \div \sqrt{(-5a)^2}$ 일 때, $10A$ 의 값을 구하여라.

 답: $10A =$ _____

20. $x > 0, y < 0$ 일 때, 다음 식을 간단히 한 것 중 옳은 것을 모두 고르면?

- ㉠

$\sqrt{(x-y)^2} = x-y$
- ㉡

$\sqrt{x^2} + \sqrt{y^2} + \sqrt{(y-x)^2} = 2x$
- ㉢

$\sqrt{x^2} - \sqrt{y^2} - \sqrt{(x-y)^2} = 2y$

- ① ㉠
- ② ㉡
- ③ ㉢
- ④ ㉠,㉡
- ⑤ ㉠,㉢