

1. 다음 중 $\sqrt{18} + 2\sqrt{2} - \frac{2}{\sqrt{2}}$ 을 바르게 계산한 것은?

- ① $\sqrt{2}$ ② $2\sqrt{2}$ ③ $3\sqrt{2}$ ④ $4\sqrt{2}$ ⑤ $5\sqrt{2}$

해설

$$\begin{aligned}(\text{준식}) &= 3\sqrt{2} + 2\sqrt{2} - \frac{2 \times \sqrt{2}}{\sqrt{2} \times \sqrt{2}} \\&= 5\sqrt{2} - \sqrt{2} \\&= 4\sqrt{2}\end{aligned}$$

2. 다음 식 $\sqrt{3}(\sqrt{54} - \sqrt{6}) + \frac{\sqrt{2}}{3} \div \sqrt{8}$ 을 간단히 한 것을 고르면?

- ① $2\sqrt{2} + \frac{1}{2}$ ② $3\sqrt{2} + \frac{1}{3}$ ③ $4\sqrt{2} + \frac{1}{4}$
④ $5\sqrt{2} + \frac{1}{5}$ ⑤ $6\sqrt{2} + \frac{1}{6}$

해설

$$\begin{aligned}(\text{준식}) &= \sqrt{162} - \sqrt{18} + \frac{\sqrt{2}}{3} \times \frac{1}{\sqrt{8}} \\&= 9\sqrt{2} - 3\sqrt{2} + \frac{1}{6} \\&= 6\sqrt{2} + \frac{1}{6}\end{aligned}$$

3. $a = \sqrt{3}, b = \sqrt{7}$ 일 때, $\frac{b}{a} \times \frac{a}{b}$ 의 값은?

- ① 1 ② $3\sqrt{7}$ ③ 4 ④ 21 ⑤ 49

해설

$$\begin{aligned}\frac{b}{a} &= \frac{\sqrt{7}}{\sqrt{3}} = \frac{\sqrt{7} \times \sqrt{3}}{\sqrt{3} \times \sqrt{3}} = \frac{\sqrt{21}}{3} \\ \frac{a}{b} &= \frac{\sqrt{3}}{\sqrt{7}} = \frac{\sqrt{3} \times \sqrt{7}}{\sqrt{7} \times \sqrt{7}} = \frac{\sqrt{21}}{7} \\ \therefore \frac{b}{a} \times \frac{a}{b} &= \frac{\sqrt{21}}{3} \times \frac{\sqrt{21}}{7} = \frac{\sqrt{21^2}}{21} = 1\end{aligned}$$

4. $\sqrt{15} \times \sqrt{20} = a\sqrt{3}$ 일 때, a 의 값은?

- ① 8 ② 10 ③ 12 ④ 15 ⑤ 18

해설

$$\begin{aligned}\sqrt{15} \times \sqrt{20} &= \sqrt{3 \times 5} \times \sqrt{2^2 \times 5} \\&= \sqrt{2^2 \times 3 \times 5^2} \\&= 10\sqrt{3}\end{aligned}$$

$$\therefore a = 10$$

5. 다음 중 계산 결과가 옳은 것의 개수는?

Ⓐ $2\sqrt{3} \div \sqrt{6} = \sqrt{2}$	Ⓑ $5\sqrt{2} \div \sqrt{5} = 5$
Ⓒ $\frac{9\sqrt{15}}{3\sqrt{15}} = \sqrt{3}$	Ⓓ $\frac{\sqrt{21}}{\sqrt{3}} = \sqrt{7}$
Ⓔ $8\sqrt{7} \div \sqrt{2} = 4\sqrt{2}$	

- ① 1개 ⓒ 2개 ③ 3개 ④ 4개 ⑤ 5개

해설

Ⓐ $2\sqrt{3} \div \sqrt{6} = \sqrt{2}$
Ⓑ $5\sqrt{2} \div \sqrt{5} = \sqrt{10}$
Ⓒ $\frac{9\sqrt{15}}{3\sqrt{15}} = 3$
Ⓓ $\frac{\sqrt{21}}{\sqrt{3}} = \sqrt{7}$
Ⓔ $8\sqrt{7} \div \sqrt{2} = 4\sqrt{14}$ 이므로
옳은 것은 ⓒ, ⓔ 두 개이다.

6. $\frac{12\sqrt{a}}{\sqrt{12}}$ 의 분모를 유리화하였더니 $2\sqrt{6}$ 이 되었다. 이 때, 자연수 $\frac{1}{\sqrt{a}}$ 의 값은?

- ① $\frac{\sqrt{2}}{4}$ ② $\frac{\sqrt{2}}{3}$ ③ $\frac{\sqrt{2}}{2}$ ④ $\sqrt{2}$ ⑤ $2\sqrt{2}$

해설

$$\frac{12\sqrt{a}}{\sqrt{12}} = \frac{12\sqrt{a}}{2\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{12\sqrt{3a}}{6} = 2\sqrt{3a} = 2\sqrt{6}$$

$$3a = 6^{\diamond} \text{으로 } a = 2$$

$$\therefore \frac{1}{\sqrt{a}} = \frac{1}{\sqrt{2}} = \frac{1 \times \sqrt{2}}{\sqrt{2} \times \sqrt{2}} = \frac{\sqrt{2}}{2}$$

7. $2\sqrt{6}\left(\frac{1}{\sqrt{3}} - \sqrt{6}\right) - \frac{a}{\sqrt{2}}(4\sqrt{2} - 2)$ 가 유리수가 되도록 유리수 a 의 값을 정하면?

- ① -1 ② -2 ③ -3 ④ -4 ⑤ -5

해설

$$\begin{aligned} & 2\sqrt{6} \times \frac{1}{\sqrt{3}} - 2\sqrt{6} \times \sqrt{6} - \frac{a}{\sqrt{2}} \times 4\sqrt{2} + \frac{a}{\sqrt{2}} \times 2 \\ &= 2\sqrt{2} - 12 - 4a + a\sqrt{2} \\ &= \sqrt{2}(2+a) - 12 - 4a \end{aligned}$$

유리수가 되기 위해서 $a+2=0$

$$\therefore a = -2$$

8. $\sqrt{6} \times \sqrt{3} \div \sqrt{12}$ 을 간단히 한 것은?

- ① $\sqrt{2}$ ② $2\sqrt{2}$ ③ $3\sqrt{2}$ ④ $\frac{\sqrt{6}}{2}$ ⑤ $2\sqrt{2}$

해설

$$\sqrt{6} \times \frac{\sqrt{3}}{\sqrt{12}} = \sqrt{\frac{6 \times 3}{12}} = \sqrt{\frac{18}{12}} = \sqrt{\frac{3}{2}} = \frac{\sqrt{6}}{2}$$

9. 다음 중 옳지 않은 것을 모두 고른 것은?

- Ⓐ $\sqrt{\frac{1}{3}} \sqrt{\frac{3}{4}} = \sqrt{\frac{1}{4}} = \frac{1}{2}$
Ⓑ $-\sqrt{60} \times \sqrt{\frac{2}{3}} = -\sqrt{60 \times \frac{2}{3}} = -\sqrt{40} = -2\sqrt{10}$
Ⓒ $\sqrt{3} \times \sqrt{12} = 6$
Ⓓ $\sqrt{0.1} \times \sqrt{0.9} = \sqrt{0.09} = 0.03$
Ⓔ $3\sqrt{5} \times 2\sqrt{7} = 6\sqrt{35}$

① Ⓐ, Ⓑ ② Ⓑ, Ⓒ ③ Ⓑ, Ⓓ ④ Ⓑ, Ⓒ ⑤ Ⓒ, Ⓓ

해설

$$\begin{aligned}\text{Ⓑ } -\sqrt{60} \times \sqrt{\frac{2}{3}} &= -\sqrt{60 \times \frac{2}{3}} \\ &= -\sqrt{40} = -2\sqrt{10}\end{aligned}$$

$$\text{Ⓓ } \sqrt{0.1} \times \sqrt{0.9} = \sqrt{0.09} = 0.3$$

10. $\sqrt{0.24} \div \sqrt{0.06} \div \sqrt{0.04}$ 를 간단히 하면?

- ① 6 ② 8 ③ 10 ④ 12 ⑤ 14

해설

$$\sqrt{\frac{24}{100}} \times \sqrt{\frac{100}{6}} \times \sqrt{\frac{100}{4}} = \sqrt{100} = 10$$

11. $\sqrt{5} = a$, $\sqrt{7} = b$ 라 할 때, $\sqrt{0.014}$ 를 a, b 를 사용하여 나타내면?

- ① $\frac{ab}{100}$ ② $\frac{ab}{50}$ ③ ab ④ $2ab$ ⑤ $4ab$

해설

$$\sqrt{0.014} = \sqrt{\frac{140}{10000}} = \frac{\sqrt{2^2 \times 5 \times 7}}{100} = \frac{2}{100} \times \sqrt{5} \times \sqrt{7} = \frac{1}{50}ab$$

12. $a > 0, b > 0$ 일 때, 다음 중 옳지 않은 것은?

$$\begin{array}{ll} \textcircled{1} \quad \frac{b}{\sqrt{a}} = \frac{b\sqrt{a}}{a} & \textcircled{2} \quad \frac{\sqrt{b}}{c\sqrt{a}} = \frac{\sqrt{ab}}{ac} \\ \textcircled{4} \quad \frac{\sqrt{b}}{\sqrt{a}} = \frac{\sqrt{ab}}{a} & \textcircled{5} \quad \frac{b}{c\sqrt{a}} = \frac{b\sqrt{a}}{ac} \end{array}$$

해설

$$\textcircled{3} \quad \sqrt{\frac{a}{b}} \times \frac{\sqrt{b}}{\sqrt{b}} = \frac{\sqrt{ab}}{b}$$

13. $a = \sqrt{2}$ 일 때, $b = 2a - \frac{3}{a}$ 이면 b 는 a 의 몇 배인가?

- ① 2 배 ② $\sqrt{2}$ 배 ③ $\frac{3}{2}$ 배
④ $\frac{1}{2}$ 배 ⑤ 3 배

해설

$$\begin{aligned} b &= 2a - \frac{3}{a} = 2\sqrt{2} - \frac{3}{\sqrt{2}} \\ &= 2\sqrt{2} - \frac{3\sqrt{2}}{2} = \left(2 - \frac{3}{2}\right)\sqrt{2} = \frac{1}{2}a \end{aligned}$$

14. $\sqrt{3}(\sqrt{10} - \sqrt{15}) \div \sqrt{5}$ 를 계산하면?

- ① $\sqrt{6} - 3$ ② $6 - \sqrt{3}$ ③ $\sqrt{6} - \sqrt{3}$
④ $\sqrt{6} + 3$ ⑤ $\sqrt{6} + \sqrt{3}$

해설

$$\begin{aligned}(\text{준식}) &= (\sqrt{3}\sqrt{10} - \sqrt{15}\sqrt{3}) \div \sqrt{5} \\&= (\sqrt{30} - \sqrt{45}) \div \sqrt{5} \\&= \sqrt{6} - \sqrt{9} \\&= \sqrt{6} - 3\end{aligned}$$

15. $f(x) = \sqrt{x+2} - \sqrt{x+1}$ 일 때, $f(0) + f(1) + f(2) + \dots + f(99) + f(100)$ 의 값을 구하면?

- ① -1 ② $\sqrt{101} - 1$ ③ $\sqrt{102} - 1$
④ $\sqrt{102} - \sqrt{101}$ ⑤ $\sqrt{102}$

해설

$$\begin{aligned}f(0) &= \sqrt{2} - \sqrt{1} = -1 + \sqrt{2} \\f(1) &= \sqrt{3} - \sqrt{2} = -\sqrt{2} + \sqrt{3} \\f(2) &= \sqrt{4} - \sqrt{3} = -\sqrt{3} + \sqrt{4} \dots \\f(99) &= \sqrt{101} - \sqrt{100} = -\sqrt{100} + \sqrt{101} \\f(100) &= \sqrt{102} - \sqrt{101} = -\sqrt{101} + \sqrt{102} \\\therefore f(0) + f(1) + f(2) + \dots + f(99) + f(100) &= -1 + \sqrt{2} - \sqrt{2} + \sqrt{3} - \sqrt{3} + \sqrt{4} + \dots - \sqrt{100} + \sqrt{101} - \sqrt{101} + \sqrt{102} \\&= -1 + (\sqrt{2} - \sqrt{2}) + (\sqrt{3} - \sqrt{3}) + (\sqrt{4} + \dots - \sqrt{100}) + (\sqrt{101} - \sqrt{101}) + \sqrt{102} \\&= -1 + (0) + (0) + (0) + \sqrt{102} \\&= -1 + \sqrt{102}\end{aligned}$$