

1. $\sqrt{96} + \frac{\sqrt{3}(\sqrt{2} - \sqrt{6})}{\sqrt{2}} - \frac{\sqrt{6} - 1}{\sqrt{2}} \div \frac{2\sqrt{2}}{\sqrt{3}}$ 를 간단히 하면?

- ① $4\sqrt{6} - \frac{5}{4}\sqrt{3} - \frac{3}{4}\sqrt{2} - 3$ ② $4\sqrt{6} + \frac{5}{4}\sqrt{3} - \frac{3}{4}\sqrt{2} - 3$
③ $4\sqrt{6} - \frac{5}{4}\sqrt{3} + \frac{3}{4}\sqrt{2} - 3$ ④ $4\sqrt{6} - \frac{5}{4}\sqrt{3} - \frac{3}{4}\sqrt{2} + 3$
⑤ $4\sqrt{6} + \frac{5}{4}\sqrt{3} + \frac{3}{4}\sqrt{2} - 3$

2. 다음 보기 중에서 옳지 않은 것은?

- ① $\sqrt{8} - \sqrt{18} + \sqrt{32} = 3\sqrt{2}$
- ② $\sqrt{27} - \sqrt{48} + \sqrt{75} = 4\sqrt{3}$
- ③ $-\frac{2}{\sqrt{2}} + \frac{32}{\sqrt{32}} = 4\sqrt{2}$
- ④ $\sqrt{5} + \sqrt{125} - \sqrt{32} + 3\sqrt{2} = 6\sqrt{5} - \sqrt{2}$
- ⑤ $\sqrt{12} + 3\sqrt{3} - \sqrt{7} + \sqrt{63} = 5\sqrt{3} + 2\sqrt{7}$

3. $(\sqrt{5} + 2\sqrt{3})(2\sqrt{5} - 3\sqrt{3})$ 을 계산하면?

- | | |
|----------------------------------|----------------------------------|
| ① $-8 - 15\sqrt{3} - 4\sqrt{15}$ | ② $-8 - 15\sqrt{3} + 4\sqrt{15}$ |
| ③ $-8 + \sqrt{15}$ | ④ $8 - 15\sqrt{3}$ |
| ⑤ $8 - 15\sqrt{3} + 4\sqrt{15}$ | |

4. $0 < x < 2$ 일 때,
 $\sqrt{(-x)^2} - \sqrt{(x-2)^2} + \sqrt{(2-x)^2}$ 을 간단히 하면?

- ① x ② $4 - x$ ③ $x + 4$
④ $3x + 4$ ⑤ $4 - 3x$

5. $-1 < a < 2$ 일 때, $\sqrt{(a+1)^2} + \sqrt{(a-2)^2} + a - 3$ 을 간단히 하면?

- | | | |
|-----------------------------|------------------------------|-------------------------|
| <p>① a</p> | <p>② $3a - 4$</p> | <p>③ 0</p> |
| <p>④ $a - 6$</p> | <p>⑤ $3a + 1$</p> | |

6. 두 실수 a, b 에 대하여 $a > 0, b < 0$ 일 때, $\sqrt{a^2 - |b|} + \sqrt{(a - b)^2}$ 을 간단히 하면?

① 0

② $2a$

③ $2b$

④ $a - b$

⑤ $2a - 2b$